

Post-harvest impact of cassava brown streak disease in four countries in eastern Africa

RODERICK J. HILLOCKS and MIDATHARAHALLY N. MARUTHI

Cassava brown streak disease (CBSD) is endemic to the coast of East Africa and the disease continues to spread to new areas in the Great Lakes region. In both these areas CBSD leaf symptoms occur at high incidences. However, it is the associated symptom of necrosis in the starch-bearing tissues that renders the root unfit for human consumption. Where root necrosis occurs, the economic viability of processing into flour is adversely affected. Surveys undertaken in Tanzania, Kenya, Uganda, and Malawi showed that CBSD leaf symptoms were present at high incidences but root necrosis incidence was lower than would be expected. It appears that a form of tolerance to CBSD occurs in which the plants are susceptible to infection by the virus but are less affected by the root symptom. Farmers practise selection for cassava varieties less prone to CBSD root necrosis leading to dependence on a decreasing number of varieties.

Keywords: cassava, CBSD, root necrosis, yield loss

WITHIN THE LAST TWO DECADES cassava brown streak disease (CBSD) has spread from the East African coast, where it has been endemic for more than 70 years (Hillocks and Jennings, 2003), to affect most of the countries of the Great Lakes region. During and prior to the 1990s, CBSD had been found at high incidences in cassava fields from the Kenyan littoral down to the Zambezi River in Mozambique, including the Lake shore region of Malawi. At the time, the disease was confined to coastal zones and cassava fields below 1,000 m above sea level. However, by around 2004, leaf symptoms resembling those of CBSD were being increasingly seen in Uganda (Alicai et al., 2007) and, as the symptoms began to spread across Uganda and into the neighbouring countries of Kenya, Tanzania, and later into Burundi, Rwanda, and the Democratic Republic of Congo, the cassava brown streak viruses (CBSVs) were confirmed as the cause (Patil et al., 2014). The viruses can be introduced into previously unaffected fields through the planting of infected cassava cuttings (the normal method of propagation) and within fields and between neighbouring fields by an insect vector, the whitefly (*Bemisia tabaci*) (Maruthi et al., 2005).

Rory Hillocks (r.j.hillocks@gre.ac.uk) is an agriculturalist at the Natural Resources Institute, Chatham, UK, with 30 years of experience in crop management for African smallholders; Midatharahally Maruthi (m.n.maruthi@gre.ac.uk) is a plant virologist at the Natural Resources Institute, Chatham, UK, acknowledged for his research on virus transmission on and host-plant resistance to CBSD.

© Practical Action Publishing, 2015, www.practicalactionpublishing.org
<http://dx.doi.org/10.3362/2046-1887.2015.008>, ISSN: 2046-1879 (print) 2046-1887 (online)




Figure 1 Scoring system used to assess severity of root necrosis

Importance of CBSD

The 'syndrome' of CBSD includes symptoms on the leaves, stem, and roots but it is the symptom of root necrosis which is most damaging and can cause total crop loss in individual plants. Infection by the virus may cause a decrease in root weight, the extent of which depends on cassava variety. The main source of crop loss is, however, the symptom of root necrosis that renders the root partly or completely unsuitable for human consumption (Hillocks and Jennings, 2003) (Figure 1). As cassava becomes increasingly used in Africa as a source of value added products and as a source of income for smallholders, CBSD poses a potential constraint to root supply for domestic and commercial processing.

Cassava processing

In much of sub-Saharan Africa between the two tropics, cassava is second to maize in importance as a starch staple and in some parts it is the primary source of carbohydrate. Even where maize is the dominant staple, cassava is grown in addition and serves a crucial role as an insurance against failure of the cereal crop. The sweet varieties of cassava can be eaten fresh but surplus production and the bitter varieties are processed into flour for cooking in the home or for sale. The traditional method of processing is to peel the roots, cut them into smaller pieces, sometimes ferment

them and/or dry them for preservation. There is a thriving local market for this dried product which is milled into flour with a fermented flavour.

Cassava can also be processed into high quality cassava flour (HQCF) that can substitute for wheat or maize flour in a number of bakery and industrial products. Partial or complete substitution of cassava flour for wheat flour in bakery products takes advantage of a locally grown crop and could significantly decrease the reliance on imported wheat in a number of African countries.

Effect of CBSD on root quality

Costs of production and aggregation are high for cassava in Africa due to low yields, its high volume to weight ratio, and poor rural road and transport networks. The profitability of community-based and larger commercial HQCF processing depends on efficient conversion of fresh root to flour. CBSD root necrosis has several effects on processing efficiency. First, roots may become scarce in areas where CBSD is prevalent, resulting in higher prices. Second, if diseased roots or portions of the root have to be discarded, the conversion ratio decreases below the optimum of 3.5 or 4:1. Third, the labour required to peel and chip the roots before grating is increased if diseased portions have to be excised. As cassava processing is increasingly seen as an income-generating option for households and small enterprises, it is important to understand the impact of CBSD on cassava production, consumption, and processing.

Impact of CBSD on production and processing

Recent research conducted in villages in Kenya, Tanzania, Uganda, and Malawi confirmed the widespread occurrence of CBSD leaf symptoms at high incidences (above 50 per cent). Surprisingly, this was not associated with a similarly high incidence of root necrosis. Severe root necrosis was found only in a few varieties usually representing a small proportion of the cassava crop. Nevertheless, moderately severe necrosis (grade 3 or above on a 1–5 scale) was found in 6–13 per cent of the cassava roots examined (Figure 2). Such production losses may not be significant in areas where root yields are high, but might decrease food security in areas such as the Lake Zone of Tanzania where in Sengerema District yields are only 8 kg ha⁻¹. CBSD root necrosis causes losses for household consumption when the necrotic portions have to be cut out before food can be prepared. Furthermore, roots affected by CBSD cannot be sold to processors. HQCF cannot be made from roots showing CBSD necrosis severity score of three or above due to decreased pH and starch content and a dark colour. Exact losses are difficult to estimate as they differ for household consumption and for processing. When cassava is scarce households may remove necrotic areas and sometimes use roots with a severity score of three. For a single household the proportion of their cassava affected by CBSD necrosis depends on the varieties they grow. For commercial processing, roots with a score of three or above are not acceptable and so the value shown in Figure 2 for percentage of roots with




Figure 2 Incidence of CBSD leaf symptoms and percentage of harvested roots showing CBSD necrosis symptom severity of grade 3 or above (1–5 scale where grade 1 = no disease) (TZ = Tanzania)

necrosis score above three is also the loss estimate. With around 36 million tonnes of cassava produced annually in the survey area, 8 per cent loss would be just under 3 million tonnes, valued at US\$750 m, based on \$250 per metric tonne.

Impact of CBSD on processing enterprises

Households view CBSD as a problem and estimate losses to the disease at 10–25 per cent. Both households and processors may have to use roots affected by CBSD when cassava is scarce and then must cut out the affected portions. Research has shown that the amount of tissue lost by cutting out diseased portions ranges from 10–60 per cent for roots showing CBSD severity grades 2–5 (Figure 3) (Abayomi et al., in press). However, a type of tolerance to CBSD occurs in the general cassava gene pool in which a variety may be fully susceptible to infection by the viruses and show typical leaf symptoms but has a low propensity to develop root necrosis. Communities in areas with a high incidence of CBSD replace the root necrosis susceptible types with the tolerant ones. As a consequence processors are still able to obtain sufficient roots free of CBSD necrosis and the impact of CBSD on the profitability of processing enterprises is less than would otherwise be the case. The main impact of CBSD in these areas has been to decrease the number of cassava varieties being grown. This loss of germplasm is particularly marked in eastern Uganda and the Lake shore region of Malawi, where there is almost total reliance on a single variety (Table 1). This poses a food security risk and a risk to processing enterprises, should those varieties later succumb to CBSD or another pest or disease epidemic.




Figure 3 Percentage loss of starch-bearing tissue when portions affected by CBSD root necrosis grades 1–5 are removed before processing

Source: Data from a community processing enterprise in Tanzania

Table 1 Number of different cassava varieties found during disease surveys conducted in selected villages in four countries in sub-Saharan Africa

Location	No of varieties	% of crop derived from 3 most common varieties
Kenya (Migori County)	14	57 (42% Weite)
Tanzania (Lake Zone)		
Village 1	5	82 (52% Mwanaminzi)
Village 2	4	94 (60% Ngarabuto)
Village 3	6	78 (47% Lufaili)
Tanzania (Southern Zone)		
Village 1	6	69 (29% Musa Saidi)
Village 2	7	77 (34% Kigoma Red)
Village 3	6	81 (38% Kigoma Red)
Malawi (Lake shore)		
Village 1	3	92 (41% Twenty:Twenty)
Village 2	3	91 (65% Twenty:Twenty)
Village 3	3	84 (78% Twenty:Twenty)
Uganda (Eastern)		
Village 1	3	91 (83% Migyera)
Village 2	2	99 (57% Migyera)
Village 3	2	100 (93% Migyera)

The adverse impact of CBSD is much greater when the disease first arrives in an area before farmers identify varieties that display tolerance to CBSD, manifested as a decreased propensity to develop root necrosis.

Long-term solution to CBSD?

As CBSD continues to spread in eastern and central Africa the disease must be managed to decrease its impact on household food security and income from processing. It seems that farmers themselves have recognized that some varieties show tolerance to CBSD and that they are less prone to root necrosis. However, these varieties are fully susceptible to infection by the viruses and their widespread propagation maintains virus inoculum at high levels. The long-term solution is to develop varieties with true resistance to CBSD in which virus replication is retarded and which show a low incidence of both leaf and root symptoms. This type of resistance is seen in the varieties Namikonga and Migyera and it will be important to fully exploit this resistance by identifying more sources of resistance and to understand the underlying mechanisms and genetics.

Conclusions

CBSD is widely distributed in eastern and central Africa and occurs at high incidences in cassava fields in the principal cassava-growing areas of at least seven countries.

The leaf symptoms are the most visible manifestation of CBSD but the most damaging symptom is root necrosis which can render the roots unfit for processing and in severe cases unfit for human consumption.

When CBSD root necrosis occurs, the profitability of cassava processing enterprises is adversely affected by the additional time taken to cut out affected portions of the root and by a decreased root to flour conversion ratio due to the resulting tissue loss.

Roots with a CBSD necrosis severity grade of 3 or above are unsuitable for processing into HQCF due to the dark colour, decrease in pH, and starch content.

Losses to CBSD are greatest in areas more recently affected by the disease but as the epidemic establishes, households begin to discard those varieties that are most susceptible to root necrosis in favour of those that are less prone to necrosis.

Because of the widespread availability of cassava varieties that are less prone to root necrosis, although fully susceptible to infection and which show typical leaf symptoms, the adverse impact of CBSD on processing enterprises is less than might otherwise be the case.

Many useful cassava varieties are being discarded due to their susceptibility to CBSD root necrosis, particularly in Eastern Uganda and the Lake shore region of Malawi, where communities for which cassava is the primary food staple are at risk of food insecurity due to dependence on only one or two varieties.

References

Abayomi, L.A., Hillocks, R.J., Maruthi, M.N. and Westby, A. (in press) 'Impact of cassava brown streak disease on cassava processing and the value chain in Eastern Africa', *Food Science Technology* (submitted May 2015).

Alicai, T., Omongo, C.A., Maruthi, M.N., Hillocks, R.J., Baguma, Y., Kawuki, R., Bua, A., Otim-Nape, G.W. and Colvin, J. (2007) 'Re-emergence of cassava brown streak disease in Uganda', *Plant Disease* 91: 24–9 <<http://dx.doi.org/10.1094/PD-91-0024>>.

Hillocks, R.J. and Jennings, D.L. (2003) 'Cassava brown streak disease: a review of present knowledge and research needs', *International Journal of Pest Management* 49: 225–34 <<http://dx.doi.org/10.1080/0967087031000101061>>.

Maruthi, M.N., Hillocks, R.J., Mtunda, K., Raya, M.D., Muhanna, M., Kiozia, H., Rekha, A.R. and Colvin, J. (2005) 'Transmission of *Cassava brown streak virus* by *Bemisia tabaci* (Gennadius)', *Journal of Phytopathology* 153: 307–12 <<http://dx.doi.org/10.1111/j.1439-0434.2005.00974.x>>.

Patil, B.L., Legg, J.P., Kanju, E. and Fauquet, C.M. (2014) 'Cassava brown streak disease: a threat to food security in Africa', *Journal of General Virology* [online] <<http://dx.doi.org/10.1099/jgv.0.000014>>.