

GUNDRUK

PICKLED LEAFY VEGETABLE

Gundruk is particularly popular in Nepal. The annual production of gundruk in Nepal is estimated at 2,000 tons and most of the production is carried out at the household level. Gundruk is obtained by fermenting and drying leafy vegetables (saag) to produce a sour brownish black product.

It is served as a side dish with the main meal and is also used as an appetiser and can be made into a soup.

Gundruk is an important source of minerals particularly during the off-season in rural areas when the diet consists of mostly starchy tubers and maize which tend to be low in minerals.



Figure 1: Preparing leafy vegetables. Photo: Practical Action Nepal.



Raw material preparation

In the months of October and November, during the harvest of the first broad mustard, radish, spinach and cauliflower leaves, large quantities of leaves accumulate - much more than can be consumed fresh.

Processing

These leaves are allowed to wilt for one or two days and then shredded with a knife or sickle.

The shredded leaves are tightly packed in an earthenware pot and warm water (at about 30°C) is added to cover all the leaves. The pot is then kept in a warm place. After five to seven days, a mild acidic taste indicates the end of fermentation and the *gundruk* is removed and dried, traditionally by the sun.

This process is similar to sauerkraut production except that no salt is added to the shredded leaves before the start of *gundruk* fermentation. The ambient temperature at the time of fermentation should be about 18°C.



Figure 2: The leaves are packed into earthenware pots. Photo: Practical Action Nepal.

Pediococcus and *Lactobacillus* species are the predominant micro-organisms during *gundruk* fermentation. During fermentation, the pH drops slowly to a final value of 4.0 and the amount of

Practical Action, The Schumacher Centre, Bourton on Dunsmore, Rugby, Warwickshire, CV23 9QZ, UK T +44 (0)1926 634400 | F +44 (0)1926 634401 | E infoserv@practicalaction.org.uk | W www.practicalaction.org




Gundruk Practical Action

acid (as lactic) increases to about 1% on the sixth day. It has been found that a disadvantage with the traditional process of *gundruk* fermentation is the loss of 90% of the carotenoids, which help to produce vitamin A, probably during sun-drying. Improved methods of drying might reduce the vitamin loss. The *Sasto* solar dryer has been developed in Nepal for use in rural areas.

Once processed the dried gundruk can be kept in airtight containers for several months.

How to make gundruk flow diagram



References and further reading

- Chiuri (The Butter Tree of Nepal) Practical Action Technical Brief
- Kawal: Fermented Green Leaves, Practical Action Technical Brief
- <u>Traditional Foods: Processing for Profit</u> by P. Fellows, Practical Action Publishing, 1997
- Fermented Fruit and Vegetables: A Global Perspective by M. Battcock & S. Azam Ali FAO, 1998
- <u>Pickles</u> a selection of Practical Action Technical Briefs

Practical Action Nepal Pandol Marga, Lazimpat

P O Box 15135, Kathmandu

Nepal

Tel: + 977 1 444 6015 / + 977 1 209 4063

Fax: + 977 1 444 5995

E-mail: info@practicalaction.org.np
Website: www.practicalaction.org/nepal

Practical Action

The Schumacher Centre Bourton-on-Dunsmore

Rugby, Warwickshire, CV23 9QZ

United Kingdom

Tel: +44 (0)1926 634400 Fax: +44 (0)1926 634401

E-mail: inforserv@practicalaction.org.uk

Website: http://practicalaction.org/practicalanswers/

Practical Action is a development charity with a difference. We know the simplest ideas can have the most profound, life-changing effect on poor people across the world. For over 40 years, we have been working closely with some of the world's poorest people - using simple technology to fight poverty and transform their lives for the better. We currently work in 15 countries in Africa, South Asia and Latin America.

